




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 7/48		A1	(11) International Publication Number: WO 97/47283
			(43) International Publication Date: 18 December 1997 (18.12.97)
(21) International Application Number: PCT/US97/10427		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GH, HU, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 13 June 1997 (13.06.97)			
(30) Priority Data: 08/664,056 13 June 1996 (13.06.96) US			
(71) Applicant: ACTIVE ORGANICS, INC. [US/US]; 11230 Grader Street, Dallas, TX 75238 (US).			
(72) Inventor: NORTON, Scott, J.; 1105 Thomridge Court, Argyle, TX 76226 (US).		Published With international search report.	
(74) Agents: GORDON, Jennifer et al.; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).			

(54) Title: COMBINATION OF ACID PROTEASE ENZYMES AND ACIDIC BUFFERS AND USES THEREOF



(57) Abstract

The invention relates to compositions which comprise at least one acid protease enzyme and an acidic buffer, the acidic buffer comprising an acid and a pharmaceutically or cosmetically acceptable carrier, which compositions are useful for treating skin conditions, diseases or disorders.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**COMBINATION OF ACID PROTEASE ENZYMES
AND ACIDIC BUFFERS AND USES THEREOF**

1. FIELD OF THE INVENTION

5 This invention relates to novel compositions which
comprise at least one acid protease enzyme and an acidic
buffer, the acidic buffer comprising an acid and a
pharmaceutically or cosmetically acceptable carrier, which
compositions are useful for treating or preventing abnormal
10 skin conditions, diseases or disorders, and/or for improving
the texture or appearance of the skin, and/or for enhancing
epidermal exfoliation, and/or for enhancing epidermal cell
renewal. The present invention is also directed to methods
for the use of the compositions for treating or preventing
15 abnormal skin conditions, diseases or disorders, and/or for
improving the texture or appearance of the skin, and/or for
enhancing epidermal exfoliation, and/or for enhancing
epidermal cell renewal.

2. BACKGROUND OF THE INVENTION

20 It is well founded that exfoliation of epidermal
layers of human skin induces an increased rate of epidermal
cell renewal (E. Phillips, 1995, U.S. Pat. No. 5,431,913;
W.P. Smith, 1994, Cosmetics and Toiletries 109:41-8). The
25 human epidermis consists of multiple layers of stratified
squamous epithelial cells in a constant state of renewal.
New cells are formed first in the basal layer, which is the
most internal membrane of the epidermis. These cells are
displaced by the production of yet newer cells and
30 subsequently are transported to the external layer of the
epidermis, the stratum corneum, where they usually are shed
(exfoliated) every two to three weeks. The general health
and appearance of human skin depends greatly upon the rate of
this process.

35 Certain situations or conditions, such as aging or
exposure to the environment, can disturb this normal process
and can lead to a generally reduced rate of cell renewal as

(VanScott and Yu, 1984, J. Am. Acad. Dermatol. 11:867-79). Because the time required for a cellular layer to migrate from the basal layer to the stratum corneum increases with the subject's age, the rate of epidermal cell renewal decreases. It has been reported that in a typical twenty year old person, the cells in the outer layers of the epidermis turn over on the average, every two weeks, while cell turnover intervals of more mature skin can be as much as twice as long (E. Phillips, 1995, U.S. Pat. No. 5,431,913).

10 The decrease in cell renewal rate due to aging can be exacerbated by environmental conditions such as exposure to solar radiation and other climatic conditions (K.E. Burke, 1990, Postgraduate Medicine 88(1):207-27). A decrease in the cell renewal rate increases the time cells in the outer

15 layers of skin are exposed to environmental conditions and may lead to further and/or cumulative damage. Certain studies suggest that a decrease in epidermal cell turnover rates is associated with an increase in intercorneocyte cohesion (VanScott and Yu, 1989, Cutis 43:222-28).

20 The cells of the epidermal layer are held together by proteinaceous components such as hemidesmosomes, desmosomes, gap junctions, glycosaminoglycans, proteoglycans, and other components present in the skin, all of which have the ability to bind to each other and to cellular components.

25 This binding together of the epidermal cells is described as "intercorneocyte cohesion", and the degree of the cohesive strength involved is affected by such factors as hydration and pH. Increased intercorneocyte cohesion results in hyperkeratinization and is characterized by thick and often

30 dry or scaly skin caused by the retention of epidermal cells. The balance that exists between cohesion and cell turnover rates is responsible for normal rejuvenation of young skin and imbalances of these factors can result in the aged, rough and unattractive look of mature epidermal surfaces. The

35 search for topically active components that balance cell renewal rates and intercorneocyte cohesion is prominent in

dermatological research efforts today (VanScott and Yu, 1984, J. Am. Acad. Dermatol. 11:867-79).

Recent advances in this research area have provided a number of topically active acidic compounds which show
5 promise in this regard (Yu et al., 1978, U.S. Pat. No. 4,105,783; Yu et al., 1982, U.S. Pat. No. 4,363,815). Long term treatments with these acidic compounds result in an increased rate of epidermal cell renewal. Those acidic components known to be epidermally active, such as low
10 molecular weight hydroxy or keto acids and esters thereof, are well documented with respect to their effectiveness as keratolytic/desquamation agents. High concentrations of these acids (e.g., salicylic and glycolic acids), as well as other acids such as trichloroacetic acid, are known for their
15 ability to cause destruction of tissue at the site of application (W.L. Epstein, 1990, in Irritant Contact Dermatitis, Jackson and Glodner, eds., Marcel Decker, Inc., New York and Basel, pp. 127-165; Remington's Pharmaceutical Sciences, A. Osol, ed., Sixteenth Ed., Mack Publishing, Inc.,
20 Easton, PA., 1980). At lower concentrations, these acids have been shown to have the ability to loosen the dead cells in the keratin-rich stratum corneum and interrupt intercorneocyte cohesion thereby facilitating desquamation; this activity is called "keratolytic activity" (Remington's
25 Pharmaceutical Sciences, A. Osol, ed., Sixteenth Ed., Mack Publishing, Inc., Easton, PA., 1980; W.P. Smith, 1994, Cosmetics and Toiletries 109:41-8).

Most keratolytics are skin irritants, including those listed above. Although some keratolytics are promoted
30 as being more efficient than others, a comparison of the therapeutic index of each of these acids shows little advantage of one over another. The "therapeutic index" is an accurate method of rating the efficiency of these components which takes into account the component's keratolytic
35 efficiency as well as the levels of irritation at a given concentration. It has been shown that even low levels of these components (greater than 4% acid concentration) cause

significant skin irritation. The use of even lower levels of these components reduces irritation but generally reduces keratolytic effectiveness. Furthermore, prolonged use of the acids reduces the efficiency of cell renewal induction (W.P. Smith, 1994, *Cosmetics and Toiletries* 109:41-8). These shortcomings point out the need for a methodology to enhance the keratolytic effectiveness of these acids at non-irritating concentrations.

The application of proteolytic enzymes in topical therapy has been used for some time for such things as scar removal from burn wounds and as an adjunct to anti-microbial therapy (G. Rodeheaver, 1975, *Am. J. Surg.* 129(5):537-544). These enzymes include those generally restricted to plant sources, such as papaya (papain), fig (ficin), and pineapple (bromelain). Cosmetic formulations containing extracts from these plants and promoted as enhancing the removal of outer epidermal layers by proteolytic action have been marketed commercially. Although the substrate specificity and spectrum of activity in relation to pH suggest that the proteolytic action of these formulations should result in keratolytic activity at the outer epidermal layers, applications of proteolytic enzymes previously used in topical therapy have not been without drawbacks.

Those enzymes (e.g., papain, bromelain, and ficin) currently used in cosmetic applications generally exhibit proteolytic activity over a broad pH range of pH 3-pH 9 (Glazer and Smith, 1971, in The Enzymes, Vol. 3, P. Boyer, ed., Academic Press, New York, pp. 501-546). It probably is the broad pH range of these plant-derived enzymes that is responsible for at least some of the problems associated with prolonged skin exposure. Human epidermal systems maintain a pH gradient within the stratified layers of the skin; the outer layers have been reported to exhibit an average pH of about 5.5 (W.P. Smith, 1994, *Cosmetics and Toiletries* 109:41-48). The pH of successive layers of the epidermis increases with depth, reaching a final pH closer to the physiological range (about pH 7.4) at the dermal layer. Because the plant-

derived enzymes mentioned above remain active across this pH gradient, there is no pH control over the activity of these plant enzymes when they are applied to the skin.

The degree of therapeutic activity derived from
5 topical application of proteolytic enzymes has been governed by the intrinsic catalytic characteristics of those enzymes. The wide range of proteolytic activity, in relation to pH, exhibited by the proteases discussed above (pH 3-9) allows for little or no control over proteolytic activity by product
10 formulation (Glazer and Smith, 1971, in The Enzymes, Vol. 3, P. Boyer, ed., Academic Press, New York, pp. 501-546). The plants from which these enzymes have been obtained are known to cause irritant contact dermatitis with symptoms including itching, edema, and blistering. It has been postulated that
15 these enzymes are the primary cause of those symptoms possibly due to excessive decreases in intercorneocyte cohesion (W.L. Epstein, 1990, in Irritant Contact Dermatitis, Jackson and Glodner, eds., Marcel Decker, Inc., New York and Basel, pp. 127-165). The enzymes currently used in cosmetic
20 applications can exhibit substrate-dependent proteolytic activity throughout all layers of the epidermis. Uncontrolled proteolysis of epidermal proteins that serve to stabilize intercorneocyte cohesion could decrease this intercorneocyte cohesion excessively and cause subsequent
25 irritation. Proteolytic attack at the basal layer of the epidermis, where cell renewal originates, could cause imbalances in the rate of epidermal cell renewal. Blistering and edema could be the result of uncontrolled proteolytic attack by this class of enzymes at the basal
30 membrane. This possibility is supported by the fact that injection of these plant enzymes into the skin of mammals is known to cause edema in the skin (Morimoto et al., 1987, Ensho 7(6):563-567). This finding has been used to produce experimental edema in dermatological research. Furthermore,
35 the symptoms of irritant contact dermatitis listed above are similar to those induced by the most effective concentrations of epidermally active acidic components currently used in

topical applications. Therefore, a more controlled approach to proteolytic activity with respect to topical applications should produce a desirable result without these side effects.

Compositions have been reported which contain a
5 proteolytic enzyme, e.g., pepsin, and several acids for the dissolution of dense necrotic formations, which formations are seen, for example, with third degree burns (Soviet Union Patent No. 439288, 1974). There is no indication that these compositions are useful in enhancing epidermal exfoliation or
10 that these compositions are active only for limited time periods.

Citation or identification of any reference in the background of this application shall not be construed as an admission that such reference is available as prior art to
15 the present invention.

3. SUMMARY OF THE INVENTION

The present invention provides novel pharmaceutical and/or cosmetic compositions for treating or preventing
20 abnormal skin conditions, diseases, or disorders, and/or for improving the texture and/or appearance of the skin, and/or for enhancing epidermal exfoliation, and/or for enhancing epidermal cell renewal. The compositions comprise at least one acid protease and an acidic buffer. For purposes of this
25 invention, the acid protease is an enzyme which exhibits peptidyl hydrolase (proteolytic) activity below the average pH of the surface of the skin and is significantly inactive at a pH greater than or equal to the average pH of the surface of the skin, which is about pH 5.5 for humans (W.P.
30 Smith, 1994, Cosmetics and Toiletries 109:41-8). For men the average pH of the surface of the skin is about pH 5.3 and for women it is about pH 6 (Ohman and Vahlquist, 1994, Acta Dermato-Venereol. 74(5):375-379). For purposes of this invention, the average pH of the surface of the skin (which
35 is about pH 5.5) includes, in addition to others, gender-specific variations, such that a pH of about 5.5 includes a range of about pH 5.3, the average pH of the surface of a

man's skin, up to about pH 6, the average pH of the surface of a woman's skin. Preferably, at a pH greater than or equal to the average pH of the surface of the skin (approximately pH 5.3-6), the acid proteases useful in this invention exhibit less than about 10% of the enzymatic activity they exhibit at their respective optimal pHs which are below the average pH of the surface of the skin.

The acid protease can be in the apoenzyme, holoenzyme, isoenzyme, or zymogen form. The acid protease component of the composition can be present in an amount of about 0.001% to about 75% by weight of the final composition, preferably about 0.1% to about 50%, more preferably about 1% to about 5%. The protease(s) has a specific activity of about 1 to about 5,000 HUT units/mg as determined by the method described in Food Chemicals CODEX, 3rd ed., (1981), pp. 496-497, National Academy Press, Washington, D.C., modified as described in Section 5.1, *infra*. Preferably the protease(s) has a specific activity of about 50 to about 3000 HUT units/mg, more preferably about 500 to about 1500 HUT units/mg.

The acidic buffer is a composition which when topically applied to the skin, temporarily lowers the pH of the surface of the skin to less than about pH 5.5 but not lower than about pH 1, preferably to between about pH 2.5 and about pH 4.5. The acidic buffer composition comprises at least one acid and a pharmaceutically or cosmetically acceptable carrier, vehicle, or excipient. The acid component of the buffer can be one or more inorganic acids, or one or more organic acids, or any combination of inorganic and/or organic acids. The acidic buffer is susceptible to neutralization to the average pH of the surface of the skin over time by natural epidermal processes, such as perspiration. The time required to neutralize the acidic buffer, and thus, inactivate of the protease, will depend on the formulation of the acidic buffer. For example, shorter time periods result if the acidic buffer contains a weak acid or a weak buffering agent to counteract the relative

alkalinity of the epidermis; longer time periods result if a stronger acid is utilized or a stronger buffering agent is employed in the acidic buffer. The acid component of the buffer can be one or more inorganic acids, or one or more
5 organic acids, or any combination of inorganic and/or organic acids and can be present in an amount of about 0.001% to about 95% by weight of the final composition, preferably about 0.01% to about 25%, more preferably about 1% to about 5%.

10 Control of the time period required for the pH of the surface of the skin to return to a pH of about 5.5 after topical application of a composition of the present invention allows for control of the activity of the protease enzyme. It is through this control of proteolytic activity that the
15 present invention overcomes the drawbacks and complications found in the prior art, such as itching, burning, blistering, etc., caused by broad pH spectrum proteolytic enzymes. The period of time it takes for the surface pH of the skin to return to about pH 5.5 is determined by a number of factors,
20 including the type of skin condition, disease, or disorder that is being treated and the sensitivity of the skin of the particular subject being treated. To avoid the drawbacks and complications found in the prior art, the period of time ideally should not exceed about 4 hours for any individual
25 application of a composition of the present invention, preferably the period of time is between from about 5 minutes to about 4 hours, more preferably between from about 30 minutes to about 2 hours, most preferably between from about 30 minutes to about one hour.

30 It is to be pointed out that the pharmaceutical compositions of the present invention are those which, when administered to the skin, render a benefit or an effect of treating or preventing an abnormal biological condition, disease, or disorder. Benefits or effects of treating or
35 preventing such abnormal condition, disease, or disorder are the reduction in severity or disappearance of the symptoms or cause of the abnormal condition, disease, or disorder. The

reduction in severity or disappearance of the abnormal condition, disease, or disorder may be either in the short-term or long-term. Such abnormal biological conditions, diseases, or disorders to be treated by administering a
5 composition of the present invention include, but are not limited to, dry skin, severe dry skin, dandruff, acne, keratoses, psoriasis, eczema, skin flakiness, pruritus, age spots, lentigines, melasmas, wrinkles (both coarse and fine, caused by intrinsic as well as extrinsic damage), warts,
10 blemished skin, hyperpigmented skin, hyperkeratotic skin, inflammatory dermatoses, age-related skin changes, skin in need of cleansers, as well as the effects of skin atrophy.

It is to be further pointed out that the cosmetic compositions of the present invention are those which, when
15 administered to the skin, improve the texture or appearance thereof or enhance epidermal exfoliation and/or epidermal cell renewal, without necessarily rendering a benefit or an effect of treating or preventing an abnormal biological condition, disease, or disorder. In this context, improving
20 the texture or appearance of the skin or enhancing epidermal exfoliation and/or epidermal cell renewal is meant to encompass providing a natural-looking and/or natural-feeling surface to the skin so as to enhance the beauty and/or smoothness of the skin from its pre-treated state, or to mask
25 unwanted symptoms of an abnormal biological condition, disease, or disorder. This can include providing a temporary moisturizing effect to the epidermis. Such abnormal biological conditions, or diseases include, but are not limited to dry skin, severe dry skin, dandruff, acne,
30 keratoses, psoriasis, eczema, skin flakiness, pruritus, age spots, lentigines, melasmas, wrinkles (both coarse and fine, caused by intrinsic as well as extrinsic damage), warts, blemished skin, hyperpigmented skin, hyperkeratotic skin, inflammatory dermatoses, psoriasis, age-related skin changes,
35 skin in need of cleansers, as well as the effects of skin atrophy.

In yet another embodiment, the invention encompasses methods for treating abnormal skin conditions, diseases, or disorders including but not limited to dry skin, severe dry skin, dandruff, acne, keratoses, psoriasis, 5 eczema, skin flakiness, pruritus, age spots, lentigines, melasmas, wrinkles (both coarse and fine, caused by intrinsic as well as extrinsic damage), warts, blemished skin, hyperpigmented skin, hyperkeratotic skin, inflammatory dermatoses, age-related skin changes, and skin in need of 10 cleansers. The method comprises topically administering to an area of a subject's skin having the condition, disease, or disorder an effective amount of a composition comprising an acid protease which is enzymatically active below about pH 5.5 and an acidic buffer which lowers the surface pH of the 15 skin to below about pH 5.5 for a period of time effective for the treatment of the condition, disease, or disorder, the acidic buffer being subject to neutralization by normal epidermal processes.

In yet another embodiment, the present invention 20 provides methods for the enhancement of epidermal exfoliation and/or epidermal cell renewal comprising topically administering to an area of a subject's skin an effective amount of a composition comprising an acid protease which is enzymatically active below about pH 5.5 and an acidic buffer 25 which lowers the surface pH of the skin to below about pH 5.5 for a period of time effective for enhancing epidermal exfoliation and/or epidermal cell renewal, the acidic buffer being subject to neutralization by normal epidermal processes.

30 In another embodiment, the invention provides methods for improving the texture or appearance of the skin comprising topically administering to an area of a subject's skin an effective amount of a composition comprising an acid protease which is enzymatically active below about pH 5.5 and 35 an acidic buffer which lowers the surface pH of the skin to below about pH 5.5 for a period of time effective to improve the texture or appearance of the skin, the acidic buffer

being subject to neutralization by normal epidermal processes.

In a still further embodiment, the invention provides methods for regulating the effects of skin atrophy comprising administering to an area of a subject's skin an effective amount of a composition comprising an acid protease which is enzymatically active below about pH 5.5 and an acidic buffer which lowers the surface pH of the skin to below about pH 5.5 for a period of time effective for regulating the effects of skin atrophy, the acidic buffer being subject to neutralization by normal epidermal processes.

In a preferred embodiment for enhancing epidermal exfoliation and/or epidermal cell renewal, the composition comprises pepsin and lactic acid. In this preferred embodiment for enhancing epidermal exfoliation and/or cell renewal, the composition comprises about 1% by weight of pepsin having a specific activity of about 1000 HUT units/mg and about 3% by weight of lactic acid. In another preferred embodiment for enhancing epidermal exfoliation and/or cell renewal, the composition comprises about 1% by weight of pepsin having a specific activity of about 1000 HUT units/mg and about 1.5% by weight of lactic acid. In yet another preferred embodiment the composition comprises 1% by weight of pepsin and 1% by weight of phosphoric acid. In another preferred embodiment the composition comprises 1% by weight of pepsin and 3% by weight of phosphoric acid. In yet another preferred embodiment the composition comprises 1% by weight of pepsin and 1% by weight of citric acid. In another preferred embodiment the composition comprises 1% by weight of pepsin and 3% by weight of citric acid.

The compositions and methods of the present invention surprisingly demonstrate pharmaceutical activity or cosmetic effects against skin disorders heretofore not achieved by acids, α -hydroxycarboxylic acids, salicylic acids, or broad pH spectrum proteases by themselves.

3.1 DEFINITIONS

As used in the present invention the following terms are intended to encompass the following:

ACIDIC BUFFER: A composition comprising an
5 acid and a pharmaceutically or
cosmetically acceptable
carrier, vehicle, or excipient
which when topically applied to
10 the skin lowers the surface pH
of the skin to below about pH
5.5 and is subject to
neutralization by natural skin
processes such that the natural
15 skin processes, over time,
return the pH of the skin's
surface to normal (which is
about pH 5.5). The acid
component of the buffer can be
20 one or more inorganic acid, or
one or more organic acid, or
any combination of inorganic
and/or organic acids.

ACID PROTEASE: An enzyme which exhibits
peptidyl hydrolase
25 (proteolytic) activity at a pH
below the average normal pH of
the skin's surface, which is
about pH 5.5, and which is
insignificantly active at a pH
30 greater than or equal to such
average normal skin surface pH,
i.e., less than about 10%
activity at about pH 5.5 or
greater as compared to peak
35 activity at pH less than about
pH 5.5. The acid protease can
be in the apoenzyme,

holoenzyme, isoenzyme, or
zymogen form.

COSMETIC:

5

A formulation to be
administered to the skin which
improves the texture or
appearance thereof, without
necessarily rendering a benefit
or an effect of treating or
preventing an abnormal
biological condition or a
disease. Such improvement
includes providing a temporary
moisturizing effect to the
mammalian epidermis.

10

15 EFFECTIVE AMOUNT:

20

An amount of composition
sufficient to significantly
induce a positive modification
in the condition to be treated,
but low enough to avoid serious
side effects. The effective
amount of the composition will
vary with the particular
condition being treated, the
age and physical condition of
the subject being treated, the
severity of the condition, the
duration of the treatment, the
nature of concurrent therapy,
the specific composition
employed, the particular
pharmaceutically-acceptable
carrier or cosmetically-
acceptable carrier utilized,
and similar factors within the
knowledge and expertise of
those skilled in the art.

25

30

35

EPIDERMAL CELL RENEWAL:

5 The process by which new skin cells are formed in the basal layer, are transported to the external layer, the stratum corneum, and subsequently are exfoliated and replaced by yet newer skin cells.

EXFOLIATION:

10 The detachment and shedding of superficial cells of an epithelium or from any tissue surface.

PHARMACEUTICAL:

15 A formulation to be administered to the skin which renders a benefit or an effect of treating or preventing an abnormal biological condition
20 or a disease.

REGULATING SKIN ATROPHY:

25 The preventing, retarding, arresting, treating, or reversing the process of atrophy in mammalian skin.

30 SKIN ATROPHY: The thinning and/or general degradation of the dermis layer of mammalian skin often characterized by a decrease in collagen and/or elastin as well as decreased number, size, and doubling potential of
35 fibroblast cells. Skin atrophy is a natural result of aging, but may be caused by either intrinsic or extrinsic factors such as natural chronoaging,

photodamage, burns or chemical damage, or by exposure to pollutants or allergens, e.g., cigarette smoke. Skin atrophy is often an undesirable side effect resulting from treatment with α -hydroxycarboxylic acids or salicylic acids.

The present invention may be understood more fully by reference to the detailed description and illustrative examples which are intended to exemplify non-limiting embodiments of the invention.

4. BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a bar graph demonstrating the effect on the rate of cell renewal with different concentrations of an acid protease (HUT Units $\times 10^6$ /g) (pepsin) with two different amounts of an acidic buffer (lactic acid). ■ 1.5% lactic acid, □ 3% lactic acid.

Figure 2 is a bar graph demonstrating the percent increases in exfoliation/cell renewal rates given by various concentrations of an acid protease (HUT Units $\times 10^6$ /g) (pepsin) over that of an acidic buffer alone (lactic acid) at concentrations of 1.5% and 3%. ■ 1.5% lactic acid, □ 3% lactic acid.

Figure 3 is a bar graph demonstrating the difference in activity of pepsin and AFP 2000, Solvay, Inc., Elkhart IN, a fungally-derived acid protease, at varying pH levels in the HUT assay. ■ AFP 2000, □ pepsin.

5. DETAILED DESCRIPTION OF THE INVENTION

5.1 COMPOSITIONS

The present invention provides novel compositions for treating or preventing abnormal skin conditions, diseases or disorders, and/or for improving the texture or appearance

of the skin, and/or for enhancing epidermal exfoliation, and/or for enhancing epidermal cell renewal. The compositions comprise at least one acid protease and an acidic buffer. Such compositions are preferably administered 5 topically, so as to minimize systemic effects or undesirable side effects. The inventor does not wish to be limited by a specific mode of action; however, the inventor believes that the compositions of the present invention accomplish the methods of the present invention by first lowering the pH of 10 the surface of the skin to a value lower than the average pH of the skin, which is about 5.5 for humans. The reduction in pH allows for the activation of the acid protease component of the composition which exhibits proteolytic/keratolytic activity during the time that the skin's surface pH is below 15 about 5.5. This time period depends upon the skin pH conditions of the individual and can be regulated by the formulation of the acidic buffer component. It is within the skill of one in the art to adjust the formulation so as to vary the time period and achieve the desired result.

20 During this time period, the acidic buffer is neutralized by natural epidermal processes, e.g., perspiration, such that the pH of the skin's surface returns to normal, which averages about pH 5.5. The increase in pH thus inactivates the acid protease and 25 proteolytic/keratolytic activity ceases, i.e., decreases by about 90%. The period of time in which the pH of the skin's surface is below about pH 5.5 and in which the protease component is active, is between from about 1 minute to about 4 hours, preferably between from about 30 minutes to about 2 30 hours, more preferably between from about 30 minutes to about 1 hour.

For purposes of this invention, the acid protease is an enzyme which exhibits significant peptidyl hydrolase (proteolytic) activity below about pH 5.5, which is the 35 average normal pH of the surface of the skin, and which is significantly inactive at a pH of about 5.5 or greater. For men the average pH of the surface of the skin is about pH

5.3 and for women it is about pH 6. For purposes of this invention, the average pH of the surface of the skin which is about pH 5.5 includes, in addition to others, gender-specific variations, such that a pH of about 5.5 includes a range of about pH 5.3, the average pH of the surface of a man's skin, up to about pH 6, the average pH of the surface of a woman's skin. The protease is significantly inactive when its activity is equal to or less than 10% activity as compared, for example, to its activity shortly after being applied to the skin as measured by the dansyl chloride test described in Section 6.1. Acid proteases useful in the present invention are those which, in an in vitro assay at pHs of about 5.5 and greater, display less than about 10% of the enzymatic activity measured in the same in vitro assay at the optimum pH (less than about pH 5.5) of the particular protease. See, e.g., Section 6.4, *infra*.

The acid protease can be in the apoenzyme, holoenzyme, isoenzyme, or zymogen form. Further, the acid proteases can be isolated from any source known to those of skill in the art, such as, for example, bacteria, fungi, tissue culture cells, and animals. The acid protease preparation can be present in an amount of about 0.001% to about 75% by weight of the final composition, preferably about 0.1% to about 50%, and most preferably about 1% to about 5%. The protease(s) will have a specific activity of about 1 to about 5000 HUT units/mg, preferably about 50 to about 3000 HUT units/mg, and most preferably about 500 to about 1500 HUT units/mg as determined by the method described in Food Chemicals CODEX, 3rd ed., (1981), pp. 496-497, National Academy Press, Washington, D.C., modified as described below.

Briefly, one first prepares the following stock solutions: HEMOGLOBIN SUBSTRATE is prepared by mixing 2 g bovine hemoglobin into 80 ml distilled water. The solution is then titrated to pH 2 by adding, for example, phosphoric acid and/or citric acid and additional distilled water is added for a total volume of 100 ml. This solution is

separated into four equal portions and each portion is titrated to the desired pH with 50% sodium hydroxide or 50% hydrochloric acid. These final solutions are then heated at 30°C for 20 minutes and are then filtered through glass wool.

5 TCA STOCK is prepared by dissolving trichloroacetic acid in distilled water for a final concentration of 5% (w/v) TCA.

Next, for each sample for which proteolytic activity is to be measured, prepare tubes labeled "B" and "T". In each tube place 4 ml of the hemoglobin solution and
10 place at 37°C such that the sample is prewarmed to 37°C. Into the "T" tube, add 100 µg enzyme solution, swirl gently and incubate at 37°C for 20 minutes. Next add 10 ml TCA stock solution to each tube. Add an equal amount of the enzyme as above added to the "T" tube to the tube marked "B" which is
15 the control for background absorbance. Centrifuge each tube and filter each sample through a syringe filter and place the filtered sample into a quartz cuvette for reading the absorbance at 280 nm. The actual absorbance is determined by subtracting the "T" sample's absorbance from the background.

20 A standard curve can be generated by measuring known quantities of protease activity.

One HUT unit of proteolytic activity is defined as that amount of enzyme that produces, in one minute under the specified conditions of the assay, a hydrolysate whose
25 absorbance at 280 nm is the same as that of a solution containing 1.10 µg per ml of tyrosine in 0.006 N hydrochloric acid. HUT units per gram are determined by the following formula:

$$\text{HUT/g} = (\text{absorbance at 280 nm} \times V) / (0.0084 \times T \times W)$$

30 where V is the final volume of the test solution, T is the reaction time in minutes, and W is the dry weight of the original enzyme sample used in the assay in grams. Protein concentration is determined by any method known in the art, such as, for example, the Bradford Assay which is described
35 in Current Protocols in Molecular Biology, Ausubel et al., (Eds.), John Wiley & Sons, Inc., New York, 1994.

Examples of such proteases include, but are not limited to, pepsin, cathepsin, human urinary acid protease, fungal proteases derived from *Neurospora oryzae*, *Mucor pusillus*, *Mucor miehei*, *Rhizopus chinensis*, or *Endothia parasitica*, bacterial proteases rhizopuspepsin, penicillopepsin, and endothiapepsin. Further, the proteases may be derived from processes involving genetic engineering processes and techniques, whether involving embryonic, mature, or induced cells and products including DNA segments, plasmids, vectors, or expression thereof, or transformed cells, or pure cultures. It should be understood that two or more acid proteases can be used in combination such that the combined amount in weight percent is within the ranges mentioned above.

The acidic buffer is a composition which when topically applied to the skin, temporarily lowers the pH of the surface of the skin to less than about pH 5.5, but not lower than about pH 1, preferably to about pH 2.5 to about pH 4.5. The acidic buffer composition comprises as least one acid and a pharmaceutically or cosmetically acceptable carrier, vehicle, or excipient. The acid component of the buffer is susceptible to neutralization to the average normal pH of the surface of the skin over time by natural epidermal processes, for example, by perspiration. The time required to neutralize the acidic buffer, and thus inactivate of the protease, will depend on the formulation of the acidic buffer. For example, shorter time periods result if the acid and/or buffering agent in the acidic buffer are weak relative to the neutralizing capacity of the epidermis; longer time periods result if a stronger acid is utilized or a stronger buffering agent is employed in the acidic buffer.

The acid component of the acidic buffer of the composition can be one or more organic acid, or one or more inorganic acid, or any combination of inorganic and/or organic acids and can be present in an amount of about 0.001% to about 95% by weight of the final composition, preferably about 0.01% to about 25%, more preferably about 1% to about

5%. The weight percent of the acid component will depend on acid strength and molarity. Moreover, the amount and strength of the acid component in the composition should be such that effective levels of proteolytic/keratolytic activity occurs but significant levels of skin irritation do not. Further, the acid component of the acidic buffer may or may not comprise an acid which has not been shown to exhibit any keratolytic activity but which merely provides an acidic environment for the activation of the protease. Examples of such acids include any molecule that can be manipulated in any composition to buffer the epidermal pH at pH ranges from about 0.1 to about 7 including, but not limited to, lactic acid, sorbic acid, phosphoric acid, citric acid, glycolic acid, malic acid, gluconic acid, pyrophosphoric acid, triphosphoric acid, polyphosphoric acid, sodium bisulfate, and potassium bisulfate. It should be understood that two or more acids can be used in combination such that the combined amount in weight percent is within the ranges mentioned above.

The acidic buffer also contains a component which is a pharmaceutically or cosmetically acceptable carrier, vehicle, or excipient. Examples of such pharmaceutically acceptable carriers, vehicles, or excipients are well known to those skilled in the art and can be found, for example, in Remington's Pharmaceutical Sciences, Eighteenth Edition, A.R. Gennaro, Ed., Mack Publishing Co., Easton, Pennsylvania, 1990. Examples of such cosmetically acceptable carriers, vehicles, or excipients are well known to those skilled in the art and can be found, for example, in CTFA International Cosmetic Ingredient Dictionary, Fourth Edition, J.M. Nikitakis, Ed., The Cosmetic, Toiletry, and Fragrance Association, Washington, D.C., 1991.

The compositions of the present invention intended for topical application may contain carrier, excipient, or vehicle ingredients such as, for example, water, acetone, ethanol, ethylene glycol, propylene glycol, butane-1,3-diol, isopropyl myristate, isopropyl palmitate, mineral oil, and

mixtures thereof to form lotions, tinctures, creams, emulsions, gels, or ointments which are non-toxic and pharmaceutically, cosmetically, or dermatologically acceptable. Additionally, moisturizers or humectants can be added to the present compositions if desired. Examples of such additional ingredients useful for pharmaceutical compositions can be found in Remington's Pharmaceutical Sciences, Eighteenth Edition, A.R.Gennaro, Ed., Mack Publishing Co., Easton, Pennsylvania, 1990. Examples of such additional ingredients useful for cosmetic compositions can be found in CTFA International Cosmetic Ingredient Dictionary, Fourth Edition, J.M. Nikitakis, Ed., The Cosmetic, Toiletry, and Fragrance Association, Washington, D.C., 1991.

In addition to these and other vehicles which will be obvious to those of ordinary skill in the art, it will be understood that the pharmaceutical and cosmetic compositions of the present invention may include other ingredients such as, for example and not by way of limitation, those that improve or eradicate age spots, keratoses, and wrinkles; analgesics; anesthetics; antiacne agents; antibacterials; antiyeast agents; antifungal agents; antiviral agents; antidandruff agents; antidermatitis agents; antipruritic agents; antiemetics; antinotion sickness agents; antiinflammatory agents; antihyperkeratolytic agents; antidryskin agents; antiperspirants; antipsoriatic agents; antiseborrheic agents; hair conditioners and hair treatment agents; antiaging and antiwrinkle agents; antiasthmatic agents and bronchodilators; sunscreen agents; antihistamine agents; skin lightening agents; depigmenting agents; vitamins; corticosteroids; tanning agents; hormones; retinoids; topical cardiovascular agents; clotrimazole; ketoconazole; miconazole; griseofulvin; hydroxyzine; diphenhydramine; pramoxine; lidocaine; procaine; mepivacaine; monobenzene; erythidocaine; procaine; mepivacaine; monobenzene; erythromycin; tetracycline; clindamycin; mecloclyline; hydroquinone; minocycline; naproxen; ibuprofen;

theophylline; cromolyn; albuterol; retinoic acid; 13-cis
retinoic acid; hydrocortisone; hydrocortisone 21-acetate;
hydro-cortisone 17-valerate; hydrocortisone 17-butyrate;
betamethasone valerate; betamethasone dipropionate;
5 triamcinolone acetonide; fluocinonide; clobetasol propionate;
benzoyl peroxide; crotamiton; propranolol; promethazine;
vitamin A palmitate; vitamin E acetate; hydrophilic
thickening agents used in pharmaceutical formulations; and
mixtures thereof. Concentrations of these ingredients will
10 vary depending upon intended use, e.g., therapeutic or
cosmetic.

Control of the time period required for the pH of
the surface of the skin to return to a normal pH of about 5.5
after topical application of a composition of the present
15 invention allows for control of the activity of the protease
enzyme. It is through this control of proteolytic activity
that the present invention overcomes the drawbacks and
complications found in the prior art, such as itching,
burning, blistering, and other undesirable side effects
20 caused by broad pH spectrum proteolytic enzymes. The
determination of the period of time it takes for the surface
of the skin to return to a normal pH of about pH 5.5 is
determined by a number of factors, including the type of skin
condition, disease, or disorder that is being treated, the
25 type of skin and its location on the body, and the
sensitivity of the skin of the particular subject being
treated. To avoid the drawbacks and complications found in
the prior art, the period of time ideally should not exceed
about 4 hours for any individual application of a composition
30 of the present invention. The time limit will depend on a
number of factors, such as the purpose of the application,
for example, day to day maintenance, removal of rough dry
skin, or therapeutic/pharmaceutical indications. Other
factors include skin type, whether its sensitive, normal,
35 insensitive, damaged, or severely damaged and the mode of
application, for example, a single application, or a
continuous wash, or the spreading of a cream or lotion.

Another factor is the specific protease or proteases used in the acid protease component because protease activity is substrate-dependent.

5

5.2 METHODS OF USE OF THE COMPOSITIONS OF THE PRESENT INVENTION

The present invention provides methods for enhancing epidermal exfoliation and/or for enhancing epidermal cell renewal. Such methods comprise topically administering to an area of a subject's skin an effective amount of a composition of the present invention, which comprises an acid protease enzyme exhibiting proteolytic activity below about pH 5.5 and insignificant activity at or above about pH 5.5 and an acidic buffer which lowers the pH of the surface of the skin to below about pH 5.5 for a period of time effective to enhance epidermal exfoliation and/or enhance epidermal cell renewal, the acidic buffer being subject to neutralization by natural epidermal processes.

The present invention also provides methods for improving the texture and/or appearance of skin. Such methods comprise administering to an area of a subject's skin an effective amount of a composition of the present invention, which comprises an acid protease enzyme exhibiting proteolytic activity below about pH 5.5 and insignificant activity at or above about pH 5.5 and an acidic buffer which lowers the pH of the surface of the skin to below about pH 5.5 for a period of time effective to improve the texture and/or appearance of skin, the acidic buffer being subject to neutralization by natural epidermal processes.

The present invention also provides methods for regulating the effects of skin atrophy. Such methods comprise administering to an area of a subject's skin an effective amount of a composition of the present invention, which comprises an acid protease enzyme exhibiting proteolytic activity below about pH 5.5 and insignificant activity at or above about pH 5.5 and an acidic buffer which lowers the pH of the surface of the skin to below pH 5.5 for

a period of time effective to regulate the effects of skin atrophy, the acidic buffer being subject to neutralization by natural epidermal processes.

The present invention also provides methods for
5 treating or preventing an abnormal skin condition, disease, or disorder. Such methods comprise administering to an area of a subject's skin an effective amount of a composition of the present invention, which comprises an acid protease enzyme exhibiting proteolytic activity below about pH 5.5 and
10 insignificant activity at or above about pH 5.5 and an acidic buffer which lowers the pH of the surface of the skin to below pH 5.5 for a period of time effective to treat or prevent an abnormal skin condition, disease, or disorder, the acidic buffer being subject to neutralization by natural
15 epidermal processes. Such condition, disease, or disorder includes, but is not limited to, dry skin, severe dry skin, dandruff, acne, keratoses, psoriasis, eczema, skin flakiness, pruritus, age spots, lentigines, melasmas, wrinkles (both coarse and fine, caused by intrinsic as well as extrinsic
20 damage), blemished skin, hyperpigmented skin, hyperkeratotic skin, inflammatory dermatoses, age-related skin changes, skin in need of cleansers, and the effects of skin atrophy.

A preferred method of administering an effective amount of a composition of the present invention for any of
25 the methods described above on an area of skin is via topical application. The amount of the acidic buffer and acid protease in the final composition and frequency of topical application to the skin can vary widely, depending upon factors such as the particular skin disorder, the severity of
30 the skin disorder, the location and/or type of the skin involved, the subject's skin sensitivity, and the degree of treatment desired. It is well within the purview of the skilled artisan to regulate dosages according to the subject's need. It is suggested as an example that topical
35 application range from about once per week to about 4 or 5 times daily, preferably from about 3 times a week to about 3 times daily, most preferably about once or twice per day.

The final composition for topical application will comprise from about 0.001% to about 95%, preferably from about 0.01% to about 25%, most preferably from about 0.01% to about 5% of the acid component. The final composition for topical
5 application will comprise from about 0.001% to about 75%, preferably from about 0.1% to about 50%, most preferably from about 1% to about 5% of the acid protease component. Another preferred mode of administering is chronic administration. "Chronic" administration, as used herein, means that the
10 period of topical application may be over the subject's lifetime, preferably for a period of at least about one month, more preferably from about three months to about twenty years, more preferably from about six months to about ten years, more preferably still from about one year to about
15 five years, thereby resulting in the treatment or prevention of the abnormal skin condition, disease, or disorder or in the improvement of the texture or appearance of the skin, or in the enhancement of epidermal exfoliation and/or epidermal cell renewal or the regulation of the effects of skin
20 atrophy.

In another embodiment of the invention, the methods described above involve administering both an effective amount of the composition comprising an acidic buffer and an acid protease and an effective amount of one or more of a
25 sunscreensing agent, an anti-inflammatory agent, an anti-oxidant/radical scavenging agent, a chelating agent, a retinoid, self-tanning material, and/or a benzofuran derivative to the skin simultaneously. As used herein, "simultaneous application" or "simultaneously" means
30 administering the agents to the skin at the same situs on the body at about the same time. Although this can be accomplished by administering the agents separately to the skin, the preferred method is to administer to the skin a composition comprising all the desired agents. The amount of
35 sunscreensing agent administered generally ranges from about 0.02 mg to about 1 mg per cm² skin. The amount of anti-inflammatory agent administered generally ranges from about

0.005 mg to about 0.5 mg, preferably from about 0.01 mg to about 0.1 mg per cm² skin. The amount of chelating agent administered generally ranges from about 0.001 mg to about 1 mg, preferably from about 0.01 mg to about 0.5 mg, more preferably from about 0.05 mg to about 0.1 mg per cm² skin. The amount of retinoid administered generally ranges from about 0.00001 mg to about 0.02 mg per cm² skin. The amount of benzofuran derivative administered generally ranges from about 0.001 mg to about 1 mg/cm² skin per application, preferably from about 0.01 to about 0.5 mg/cm² skin per application. The amount of the composition comprising an acidic buffer and an acid protease administered generally ranges from about 0.001 mg to about 1 mg per cm² skin per application, preferably from about 0.01 mg to about 0.5 mg per cm², more preferably from about 0.05 to about 0.25 mg/cm² skin per application, depending upon the severity of the condition to be treated, the sensitivity of the subject's skin, the location of the treatment site on the subject, and the efficacy of the compounds employed.

20

6. EXAMPLES

In order to more fully illustrate the nature of the invention and the manner of practicing the same, the following examples are provided, which are not to be construed as limiting the remainder of the disclosure or the scope of the invention in any way whatsoever.

The following reagents were used in the compositions and methods described in Section 6.1 through 6.4:

AFP 2000, a fungal-derived acid protease, Solvay Inc., Elkhart IN; algae extract/aloe extract, Active Organics, Inc., Dallas TX; allantoin, Sutton Laboratories, Chatham NJ; Arlacel 65, ICI Surfactants, Wilmington DE; aspartic acid, Ajinomoto USA, Inc., Teaneck NJ; BHT, Eastman Kodak, Co., Rochester NY; butylene glycol, Hoechst Celanese, Co., Charlotte NC; Carbomer, B.F. Goodrich, Co., Brecksville OH; cetearyl alcohol, Lipo Chemicals, Inc., Patterson NJ; cetyl

alcohol, Lipo Chemicals, Inc., Patterson NJ; citric acid, Roche Vitamins and Fine Chemicals, Nutley NJ; Cosmowax, Croda Chemicals, Ltd., Parsippany NJ; cyclomethicone, Dow Chemicals USA, Midland MI; dihydroxyacetone, Rona/E. Merck Industries, Hawthorne NY; dimethicone, Dow Chemicals USA, Midland MI; disodium EDTA, Ciba-Geigy Co., Greensboro NC; DL-Panthenol, Roche Vitamins and Fine Chemicals, Nutley NJ; ethoxydiglycol, Gattefosse S.A., Paris, France; gluconic acid, Sigma Chemicals, St. Louis MO; Glycereth-26, Amerchol Co., Edison NJ; glycerol stearate, Croda Chemicals, Ltd., Parsippany NJ; lactic acid USP 88%, Rita Corporation, Woodstock IL; lecithin/asiatic acid, Active Organics, Dallas TX; methyl gluceth-20, Amerchol Co., Edison NJ; methylparaben, Napp Co., Lodi NJ; octyl palmitate, ICI Surfactants, Wilmington DE; octyldodecyl neopentanoate, ICI Surfactants, Wilmington DE; PEG-75, Lipo Chemicals, Inc., Patterson NJ; PEG-40 stearate, Lipo Chemicals, Inc., Patterson NJ; Pepsin 1:15,000 NF, American Laboratories, Inc., Omaha NE; phosphoric acid 85%, Sigma Chemicals, Inc., St. Louis MO; polyacrylamide/C13-14 isoparaffin/laureth-7, Seppic, Co., Paris, France; propylene glycol, Dow Chemicals USA, Midland MI; propylparaben, Napp Co., Lodi NJ; retinyl palmitate, Roche Vitamins and Fine Chemicals, Nutley NJ; salicylic acid, Kalama Co., Seattle WA; sodium carboxyl-methylcellulose 7HXF, Aqualon, Co., Wilmington DE; sodium hyaluronate, Active Organics, Dallas TX; steareth-10, Lipo Chemicals, Inc., Patterson NJ; tocopheryl acetate, Roche Vitamins and Fine Chemicals, Nutley NJ; triethanolamine 99%, BASF Co., Mount Olive NJ; and xanthan gum, Kelco, San Diego, CA.

30

6.1 ENHANCEMENT OF SKIN EXFOLIATION

A composition of the present invention was tested at differing concentrations of the acid protease pepsin, 1:15,000 NF, American Laboratories, Inc., Omaha, NE, and the acidic buffer lactic acid on individual human volar forearms for the enhancement of skin exfoliation and cell renewal. Twenty subjects between the ages of 30 and 60 years were

selected and were required to refrain from using any products on their volar forearm, except those supplied in conjunction with the test procedure (which included a synthetic soap) for 5 days before and during the test period. None of the 5 subjects: (1) was pregnant or lactating or intended to become pregnant within three months following the test; (2) had one or more known communicable diseases; (3) was involved in any other clinical study at the time of the test; (4) suffered from psoriasis, dermatitis, and/or eczema; (5) was taking 10 medication considered likely to interfere with the test results, including retinoids, corticosteroids, or antibiotics; (6) had taken retinoids, corticosteroids, or antibiotics during the six months prior to the test; and/or (7) had used tobacco products, had sensitive skin, had a skin 15 type deemed unsuitable for the test, or had other characteristics considered likely to make them unsuitable for the test.

Each volar forearm of each subject was patched with four 2 x 3 cm adhesive bandages, to which had been applied 1- 20 2 gm/cm² of 5% ultra-pure dansyl chloride milled into petrolatum. Three of the bandages on each forearm were used as test sites and the remaining one was used as a control.

The four sites on each forearm of each subject were covered with the dansyl chloride-loaded bandages and were 25 left undisturbed for 24 hours. At the end of the 24 hour period, the bandages were removed, the sites were washed, and staining of the sites by dansyl chloride was confirmed by viewing with a long wave UV light source. The six non-control dansyl chloride stained test sites on each subject 30 received twice daily topical applications of 1-2 ml/cm² of a randomized selection of test compositions described in Table I. Upon application, the test compositions were rubbed into the skin at the test sites until the sites were no longer wet. After the dansyl chloride staining was verified, 35 the test sites and the control sites were left uncovered and were handled in the same manner except that the test sites

received the test applications and the control sites did not receive any test composition.

TABLE I

Test Composition No.	Lactic Acid (100%) % by wt.	Acid Protease ^a % by wt.	HUT units/g * 10 ⁶	Sorbic Acid preservative % by wt.	Deionized Water % by wt.
1	0.00	0.00	0.0	0.05	99.95
2	1.50	0.00	0.0	0.05	98.45
3	1.50	0.02	0.02	0.05	98.43
4	1.50	0.20	0.2	0.05	98.25
5	1.50	1.00	1.0	0.05	97.45
6	1.50	2.00	2.0	0.05	96.45
7	1.50	4.00	4.0	0.05	94.45
8	3.00	0.00	0.0	0.05	96.95
9	3.00	0.02	0.02	0.05	96.93
10	3.00	0.20	0.2	0.05	96.75
11	3.00	1.00	1.0	0.05	95.95
12	3.00	2.00	2.0	0.05	94.95
13	3.00	4.00	4.0	0.05	92.95

^a The acid protease employed was Pepsin, 1:15,000, NF, American Laboratories, Inc., Omaha NE.

Exfoliation/keratolysis of the stratum corneum was determined by visualizing the dansyl chloride stains daily under a long wave UV light source to measure stain removal. The percent increase in exfoliation/keratolysis and accompanying cell renewal of the stratum corneum was calculated by the following formula:

$$\% \text{ increase} = \frac{(\text{days for removal of untreated area} - \text{days for removal of treated area})}{\text{days for removal of untreated area}} \times 100$$

5

TABLE II

Test Composition No.	Lactic Acid Conc. (%)	Acid Protease* Conc. (%)	HUT units/g * 10 ⁴	Effect of Composition (% Increase over Control Value)	Pepsin Effect (% Increase over Lactic Acid Control Values)
1	0.00	0.00	0.0	0 (Control)	0
2	1.50	0.00	0.0	14	0 (Control)
3	1.50	0.02	0.02	14	0
4	1.50	0.20	0.2	17	21
5	1.50	1.00	1.0	25	79
6	1.50	2.00	2.0	28	100
7	1.50	4.00	4.0	33	136
8	3.00	0.00	0.0	18	0 (Control)
9	3.00	0.02	0.02	18	0
10	3.00	0.20	0.2	23	28
11	3.00	1.00	1.0	31	72
12	3.00	2.00	2.0	33	83
13	3.00	4.00	4.0	36	100

* The acid protease employed was Pepsin, 1:15,000, NF, American Laboratories, Inc., Omaha NE.

30

The data collected from the test are summarized in Table II and in Figures 1 and 2. As seen in Table II, column 5, the acidic buffer lactic acid had some positive keratolytic/cell renewal effects alone [compare test composition 2 and 8 to the control (test composition 1)] at concentrations of 1.5% and 3% by weight. These positive

keratolytic/cell renewal effects are significantly enhanced in the presence of the acid protease pepsin, (see test compositions 3-7 and 9-13). It is also apparent that the enhancements are dose dependent, i.e., stratum corneum exfoliation/cell renewal rates increase as the concentration of the acid protease increases, see column 5 of Table II. Figure 1 presents the data in column 5 graphically.

Column 6 of Table II reflects the percent increase in exfoliation/cell renewal rates given by various concentrations of the acid protease pepsin over that given by lactic acid alone at 1.5% and 3% by weight concentration. The data in column 6 is presented graphically in Figure 2. It is apparent that, although the acid protease effect is significant at both lactic acid concentrations employed, the greater percent enhancements are given by the acid protease pepsin at a lactic acid concentration of 1.5% by weight.

A low level of redness and swelling was observed on the skin of some of the test subjects, approximately 75% of the skin test sites at a 3% by weight lactic acid concentration, but very little was observed at a 1.5% by weight lactic acid concentration. Furthermore, this finding was largely independent of the presence of pepsin. Thus, significant exfoliation/cell renewal is accomplished by employing an acidic buffer at a low, non-irritating level in the presence of an appropriate concentration of an acid protease. In contrast to proteases which are active in the physiological pH range, acid proteases, as defined herein, applied to the skin maintain their proteolytic activity only until the skin's own natural processes raise the pH of the skin's surface to its normal value of approximately 5.5.

6.2 COMPOSITIONS FOR USE IN THE METHODS OF THE PRESENT INVENTION

The following Tables recite formulations of different examples of compositions of the present invention. All of the following compositions were adjusted to a pH of

about 3 to 3.2 with, for example, dilute phosphoric acid or sodium hydroxide, NaOH.

TABLE III

WASH AP-01

5

		PERCENT BY WEIGHT
	ASPARTIC ACID	1.50
	DEIONIZED WATER	82.50
10	METHYLPARABEN	0.20
	PEG 75	1.50
	DISODIUM EDTA	0.05
	ALLANTOIN	0.25
15	GLYCERETH 26	1.00
	ETHOXYDIGLYCOL	4.00
	PROPYLENE GLYCOL	4.00
	AFP 2000	5.00
20	TOTALS	100.00

25

30

35

TABLE IV

WASH AP-02

5		PERCENT BY WEIGHT
	PHOSPHORIC ACID 85%	1.20
	DEIONIZED WATER	86.80
	METHYLPARABEN	0.20
	PEG 75	1.50
10	DISODIUM EDTA	0.05
	ALLANTOIN	0.25
	GLYCERETH 26	1.00
	ETHOXYDIGLYCOL	4.00
15	PROPYLENE GLYCOL	4.00
	PEPSIN 1,000 HUT UNITS/MG	1.00
	TOTALS	100.00

20

25

30

35

Serum AP-03

TABLE V

5		PERCENT BY WEIGHT
	SALICYLIC ACID	0.50
	TRIETHANOLAMINE 99%	1.00
	LACTIC ACID USP 88%	1.20
	DEIONIZED WATER	82.70
10	SODIUM CARBOXY-METHYCELLULOSE 7HXF	0.60
	METHYLPARABEN	0.20
	PEG 75	1.50
	DISODIUM EDTA	0.05
15	ALLANTOIN	0.25
	METHYL GLUCETH-20	1.00
	GLYCERETH 26	1.00
	ETHOXYDIGLYCOL	4.00
20	PROPYLENE GLYCOL	4.00
	LECITHIN/ ASIATIC ACID	1.00
	PEPSIN 1,000 HUT UNITS/MG	1.00
25	TOTALS	100.00

30

35

TABLE VI

Serum AP-04

5		PERCENT BY WEIGHT
	GLUCONIC ACID	1.50
	DEIONIZED WATER	84.08
	METHYLPARABEN	0.02
10	SODIUM CARBOXY-METHYCELLULOSE 7HXF	0.60
	PEG 75	1.50
	DISODIUM EDTA	0.05
	ALLANTOIN	0.25
	METHYL GLUCETH-20	1.00
15	GLYCERETH 26	1.00
	ETHOXYDIGLYCOL	4.00
	PROPYLENE GLYCOL	4.00
	LECITHIN/ ASIATIC ACID	1.00
20	PEPSIN 1,000 HUT UNITS/MG	1.00
	TOTALS	100.00

25

30

35

Pepsin Lotion AP-05

TABLE VII

5		PERCENT BY WEIGHT
	DEIONIZED WATER	64.75
	XANTHAN GUM	0.10
	DISODIUM EDTA	0.10
	ALLANTOIN	0.50
10	ALGAE EXTRACT (and) ALOE EXTRACT	6.00
	METHYLPARABEN	0.20
	DL-PANTHENOL	0.50
	ETHOXYDIGYCOL	4.00
15	BUTYLENE GYCOL	5.00
	CETYL ALCOHOL	2.00
	GLYCERYL STEARATE	2.00
	OCTYL PALMITATE	4.00
20	CYCLOMETHICONE	2.00
	PROPYLPARABEN	0.10
	BHT	0.05
	POLYACRYLAMIDE (and) C 13-14 ISOPARAFFIN (and) LAURETH-7	1.50
25	SODIUM HYALURONATE	3.00
	RETINYL PALMITATE	0.10
	TOCOPHERYL ACETATE	0.10
	PEPSIN 1,000 HUT UNITS/MG	1.00
30	CITRIC ACID	3.00
	TOTALS	100.00

35

TABLE VIII

Renewal Gel AP-06

5		PERCENT BY WEIGHT
	DEIONIZED WATER	89.60
	CARBOMER	1.50
	SALICYLIC ACID	0.50
	LACTIC ACID USP 88%	1.20
10	PROPYLENE GLYCOL	6.00
	PEPSIN 1,000 HUT UNITS/MG	1.00
	METHYLPARABEN	0.15
	PROPYLPARABEN	0.05
15	TOTALS	100.00

20

TABLE IX

Renewal Gel AP-07

25		PERCENT BY WEIGHT
	DEIONIZED WATER	85.10
	POLYACRYLAMIDE (and) C13-14 ISOPARAFFIN (and) LAURETH-7	4.00
	SALICYLIC ACID	0.50
	LACTIC ACID USP 88%	1.20
30	BUTYLENE GLYCOL	8.00
	PEPSIN 1,000 HUT UNITS/MG	1.00
	METHYLPARABEN	0.15
	PROPYLPARABEN	0.05
35	TOTALS	100.00

Self-Tanning Lotion AP-08

TABLE X

5		PERCENT BY WEIGHT
	DEIONIZED WATER	48.15
	XANTHAN GUM	0.75
	PROPYLENE GLYCOL	5.00
10	METHYLPARABEN	0.20
	PROPYLPARABEN	0.10
	PEPSIN 1,000 HUT UNITS/MG	2.00
	STEARETH-10	0.70
	ARLACEL 1651	1.20
15	PEG-40 STEARATE	0.40
	COSMOWAX J2	1.00
	CETEARYL ALCOHOL	1.50
	CYCLOMETHICONE	5.00
20	DIMETHICONE	0.50
	OCTYLDODECYL NEOPENTANOATE	28.50
	DIHYDROXYACETONE	5.00
	TOTALS	100.00

25

30

35

6.3 STABILIZATION COMPOSITION

The following composition at pH 3 retains at least 90% of its acid protease activity over a period of at least three months, during which the composition was stored at 37°C.

		<u>% by weight</u>
	Cyclomethicone	5.0
	Cetyl Alcohol	1.5
10	Arlacel 165	1.5
	Dimethicone	1.0
	Sepigel 305	3.0
	Phenoxyethanol	1.0
	EDTA, disodium	0.1
15	Butylene Glycol	10.0
	Citric acid/Sodium Citrate	3.0
	Lactic acid (100%)	1.5
	Sorbic acid	0.05
	Lactose	1.0
20	Pepsin:	1.0
	1000 HUT Units/mg	
	H ₂ O	70.35

6.4 PROTEOLYTIC ACTIVITY AT VARYING pH LEVELS

Two acid protease enzymes were tested over a range of pH values to determine if they have proteolytic activity at a pH less than about pH 5.5 and whether the activity decreases to less than about 10% compared to its activity at the pH optimum. Proteolytic activity was measured by determining the proteolytic effect on hemoglobin using the HUT assay as described below. The two acid proteases were pepsin 1:15,000 NF, at about 1000 HUT units/mg, supplied by American Laboratories, Inc., Omaha NE, and AFP 2000, a fungally-derived acid protease supplied by Solvay, Inc., Elkhart IN, at about 100 HUT units/mg. The assay was performed first by preparing stock solutions HEMOGLOBIN SUBSTRATE and TCA STOCK. The hemoglobin substrate was made by dissolving 2 g of bovine hemoglobin, Sigma Chemicals,

Inc., St. Louis MO, into 80 ml distilled water. This solution was then titrated to the correct pH, either, pH 2, 3.5, 5.5 or 6. For pH values between 1 and 3, 1.8 g of phosphoric acid was dissolved for a final concentration of 100 mM and the pH was adjusted with 50% sodium hydroxide or 50% hydrochloric acid. For pH values between 3 and 6, 2 g of citric acid was dissolved for a final concentration of 100 mM and the pH was adjusted with 50% sodium hydroxide or hydrochloric acid. The solutions were brought to a final volume of 100 ml by adding distilled water. The substrates with differing pH values were then incubated at 30°C for 20 minutes and subsequently filtered through glass wool. The TCA stock solution was prepared by dissolving 5 g of trichloroacetic acid into 100 ml distilled water.

The enzyme standards were then prepared by dissolving 10 mg of pepsin in 10 ml distilled water and by dissolving 100 mg of AFP 2000 and 300 mg citric acid in 8 ml distilled water. This solution was then adjusted to a pH of about 3.5 with 6M HCl or 50% sodium hydroxide and the final volume was brought to 10 ml by the addition of distilled water.

For each of the samples, four tubes were marked, "B", "T1", "T2" and "T3", and 4 ml of the hemoglobin substrate solution was added to each tube, which were then placed at 37°C. 100 µl of the enzyme standard was added to each "T" labeled tube, i.e., T1, T2 and T3. The tubes were then incubated for 20 minutes at 37°C for pepsin and 30 minutes for AFP 2000. 10 ml of the TCA stock solution was then added to all the tubes, and 100 µl of enzyme was added to tube "B". The tubes were then centrifuged for one minute at 1000 rpm. The sample in each tube was filtered through a syringe filter into a quartz cuvette and the absorbance at 280 nm was taken. The background absorbance was deducted from each of the "T"'s absorbance and the individual "T1", "T2" and "T3" absorbencies were averaged for each enzyme at each pH tested.

Figure 3 demonstrates that both of the enzymes were optimally active at pH 2. Further, both enzymes had less

than 10% activity at pH 5.5 and greater as compared to the optimal level at pH 2. This assay clearly demonstrates that these enzymes are enzymes which can be used in the present invention. Further, by using such an assay, it may be
5 determined which additional enzymes are suitable for use in the present invention.

The invention described and claimed herein is not to be limited in scope by the specific embodiments herein disclosed since these embodiments are intended as
10 illustrations of several aspects of the invention. Any equivalent embodiments are intended to be within the scope of this invention. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the
15 foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

A number of references are cited herein, the entire disclosures of which are incorporated herein, in their entirety, by reference.

20

25

30

35

WHAT IS CLAIMED IS:

1. A composition for the treatment or prevention of an abnormal skin condition, disease, or disorder and/or for the enhancement of epidermal exfoliation and/or for improving the texture or appearance of the skin comprising:
(1) an acid protease which is enzymatically active below about pH 5.5 and which is insignificantly active at or above pH 5.5; and (2) an acidic buffer comprising an inorganic acid which temporarily lowers the surface pH of the skin to below about pH 5.5, the acidic buffer being subject to neutralization by natural epidermal processes, such that the surface pH of the skin returns to about pH 5.5.
2. The composition according to claim 1 wherein the acidic buffer further comprises a pharmaceutically or cosmetically acceptable carrier, vehicle, or excipient.
3. The composition according to claim 1 wherein the acid protease is selected from the group consisting of fungal proteases, bacterial proteases or mammalian proteases and mixtures thereof.
4. The composition according to claim 3 wherein the acid protease is selected from the group consisting of pepsin, cathepsin, human urinary acid protease, rhizopuspepsin, penicillopepsin, endothiapepsin and mixtures thereof.
5. The composition according to claim 1 wherein the inorganic acid is selected from the group consisting of phosphoric acid, pyrophosphoric acid, triphosphoric acid, polyphosphoric acid, sodium bisulfate, potassium bisulfate and mixtures thereof.
6. The composition according to claim 2 wherein the pharmaceutically or cosmetically acceptable carrier, vehicle, or excipient component of the acidic buffer is

selected from the group consisting of lotions, tinctures, creams, emulsions, gels, ointments, water, water-workable cream, polyvinyl alcohol, hydroxyethyl cellulose, cellulose, hydrophilic acrylic polymer, emollients, skin moisturizing
5 components, enzyme stabilizers, glycerol, surfactants, preservatives, hydrophilic thickening agents used in pharmaceutical formulations and mixtures thereof.

7. The composition according to claim 1 wherein
10 the acid protease is present in an amount of about 0.001% to about 75% by weight of the final composition.

8. The composition according to claim 7 wherein the acid protease is present in an amount preferably of about
15 0.1% to about 50% by weight of the final composition.

9. The composition according to claim 8 wherein the acid protease is present in an amount most preferably of about 1% to about 5% by weight of the final composition.
20

10. The composition according to claim 1 wherein the acid protease has a total specific activity of about 1 to about 5000 HUT units/mg.

25 11. The composition according to claim 10 wherein the acid protease has a total specific activity preferably of about 50 to about 3000 HUT units/mg.

12. The composition according to claim 11 wherein
30 the acid protease has a total specific activity most preferably of about 500 to about 1500 HUT units/mg.

13. The composition according to claim 1 wherein the acid component of the acidic buffer is present in an
35 amount of about 0.001% to about 95% by weight of the final composition.

14. The composition according to claim 13 wherein the acid component of the acidic buffer is present in an amount preferably of about 0.01% to about 25% by weight of the final composition.

5

15. The composition according to claim 14 wherein the acid component of the acidic buffer is present in an amount most preferably of about 1% to about 5% by weight of the final composition.

10

16. A method for enhancing epidermal exfoliation and/or for enhancing epidermal cell renewal which comprises topically administering to a subject on an area of skin an effective amount of a composition comprising (1) an acid
15 protease which is enzymatically active below about pH 5.5 and which is insignificantly active at or above pH 5.5; and (2) an acidic buffer which temporarily lowers the surface pH of the skin to below about pH 5.5 for a period of time effective
20 for enhancing epidermal exfoliation and/or epidermal cell renewal, the acidic buffer being subject to neutralization by natural epidermal processes, such that the surface pH of the skin returns to about 5.5.

17. The method according to claim 16 wherein the
25 acidic buffer further comprises a pharmaceutically or cosmetically acceptable carrier, vehicle, or excipient.

18. The method according to claim 16 wherein the acid protease is selected from the group consisting of fungal
30 proteases, bacterial proteases or mammalian proteases and mixtures thereof.

19. The method according to claim 18 wherein the acid protease is selected from the group consisting of
35 pepsin, cathepsin, human urinary acid protease, rhizopuspepsin, penicillopepsin, endothiapepsin and mixtures thereof.

20. The method according to claim 16 wherein the acid protease is pepsin.

21. The method according to claim 16 wherein the acid component of the acidic buffer is selected from the group consisting of inorganic acids, organic acids and mixtures thereof.

22. The method according to claim 21 wherein the inorganic acid is selected from the group consisting of phosphoric acid, pyrophosphoric acid, triphosphoric acid, polyphosphoric acid, sodium bisulfate, potassium bisulfate and mixtures thereof.

23. The method according to claim 21 wherein the organic acid is selected from the group consisting of lactic acid, citric acid, sorbic acid, glycolic acid, malic acid, gluconic acid and mixtures thereof.

24. The method according to claim 17 wherein the pharmaceutically or cosmetically acceptable carrier, vehicle, or excipient component of the acidic buffer is selected from the group consisting of lotions, tinctures, creams, emulsions, gels, ointments, water, water-workable cream, polyvinyl alcohol, hydroxyethyl cellulose, cellulose, hydrophilic acrylic polymer, emollients, skin moisturizing components, enzyme stabilizers, glycerol, surfactants, preservatives, hydrophilic thickening agents used in pharmaceutical formulations and mixtures thereof.

25. The method according to claim 16 wherein the acid protease is present in an amount of about 0.001% to about 75% by weight of the final composition.

26. The method according to claim 25 wherein the acid protease is present in an amount preferably of about 0.1% to about 50% by weight of the final composition.

27. The method according to claim 26 wherein the acid protease is present in an amount most preferably of about 1% to about 5% by weight of the final composition.

5 28. The method according to claim 16 wherein the acid protease has a total specific activity of about 1 to about 5000 HUT units/mg.

29. The method according to claim 28 wherein the
10 acid protease has a total specific activity preferably of about 50 to about 3000 HUT units/mg.

30. The method according to claim 29 wherein the acid protease has a total specific activity most preferably
15 of about 500 to about 1500 HUT units/mg.

31. The method according to claim 16 wherein the acid component of the acidic buffer is present in an amount of about 0.001% to about 95% by weight of the final
20 composition.

32. The method according to claim 31 wherein the acid component of the acidic buffer is present in an amount preferably of about 0.01% to about 25% by weight of the final
25 composition.

33. The method according to claim 32 wherein the acid component of the acidic buffer is present in an amount most preferably of about 1% to about 5% by weight of the
30 final composition.

34. The method according to claim 16 wherein the surface pH of the skin is below about pH 5.5 for about 5 minutes to about 4 hours.

35

35. The method according to claim 34 wherein the surface pH of the skin is below about pH 5.5 for about 30 minutes to about 2 hours.

36. The method according to claim 35 wherein the surface pH of the skin is below about pH 5.5 for about 30 minutes to about 1 hour.

5 37. A method for treating or preventing an abnormal skin condition, disease or disorder and/or for regulating the effects of skin atrophy and/or for improving the texture or appearance of the skin which comprises topically administering to a subject on an area of skin an
10 effective amount of a composition comprising (1) an acid protease which is enzymatically active below about pH 5.5 and which is insignificantly active at or above pH 5.5; and (2) an acidic buffer which temporarily lowers the surface pH of the skin to below about pH 5.5 for a period of time effective
15 for the treatment or prevention of the condition, disease, or disorder and/or for regulating the effects of skin atrophy and/or for improving the texture or appearance of the skin, the acidic buffer being subject to neutralization by natural epidermal processes, such that the surface pH of the skin
20 returns to about pH 5.5.

38. The method according to claim 37 wherein the acidic buffer further comprises a pharmaceutically or cosmetically acceptable carrier, vehicle, or excipient.
25

39. The method according to claim 37 wherein the acid protease is selected from the group consisting of fungal proteases, bacterial proteases and mammalian proteases or mixtures thereof.

30 40. The method according to claim 39 wherein the acid protease is selected from the group consisting of pepsin, cathepsin, human urinary acid protease, rhizopuspepsin, penicillopepsin, endothiapepsin and mixtures
35 thereof.

41. The method according to claim 37 wherein the acid protease is pepsin.

42. The method according to claim 37 wherein the acid component of the acidic buffer is selected from the group consisting of inorganic acids, organic acids and mixtures thereof.

5

43. The method according to claim 42 wherein the inorganic acid is selected from the group consisting of phosphoric acid, pyrophosphoric acid, triphosphoric acid, polyphosphoric acid, sodium bisulfate, potassium bisulfate
10 and mixtures thereof.

44. The method according to claim 42 wherein the organic acid is selected from the group consisting of lactic acid, citric acid, sorbic acid, glycolic acid, malic acid,
15 gluconic acid and mixtures thereof.

45. The method according to claim 38 wherein the pharmaceutically or cosmetically acceptable carrier, vehicle, or excipient component of the acidic buffer is selected from
20 the group consisting of lotions, tinctures, creams, emulsions, gels, ointments, water, water-workable cream, polyvinyl alcohol, hydroxyethyl cellulose, cellulose, hydrophilic acrylic polymer, emollients, skin moisturizing components, enzyme stabilizers, glycerol, surfactants,
25 preservatives, hydrophilic thickening agents used in pharmaceutical formulations and mixtures thereof.

46. The method according to claim 37 wherein the acid protease is present in an amount of about 0.001% to
30 about 75% by weight of the final composition.

47. The method according to claim 46 wherein the acid protease is present in an amount preferably of about 0.1% to about 50% by weight of the final composition.
35

48. The method according to claim 47 wherein the acid protease is present in an amount most preferably of about 1% to about 5% by weight of the final composition.

49. The method according to claim 37 wherein the acid protease has a total specific activity of about 1 to about 5000 HUT units/mg.

5 50. The method according to claim 49 wherein the acid protease has a total specific activity preferably of about 50 to about 3000 HUT units/mg.

3 51. The method according to claim 50 wherein the
10 acid protease has a total specific activity most preferably of about 500 to about 1500 HUT units/mg.

52. The method according to claim 37 wherein the acid component of the acidic buffer is present in an amount
15 of about 0.001% to about 95% by weight of the final composition.

53. The method according to claim 52 wherein the acid component of the acidic buffer is present in an amount
20 preferably of about 0.01% to about 25% by weight of the final composition.

54. The method according to claim 53 wherein the acid component of the acidic buffer is present in an amount
25 most preferably of about 1% to about 5% by weight of the final composition.

55. The method according to claim 37 wherein the surface pH of the skin is below about pH 5.5 for about 5
30 minutes to about 4 hours.

56. The method according to claim 55 wherein the surface pH of the skin is below about pH 5.5 for about 30 minutes to about 2 hours.

35

57. The method according to claim 56 wherein the surface pH of the skin is below about pH 5.5 for about 30 minutes to about 1 hour.

58. The method according to claim 37 wherein abnormal skin condition, disease or disorder is selected from the group consisting of dry skin, severe dry skin, dandruff, acne, keratoses, psoriasis, eczema, skin flakiness, pruritus, age spots, lentigines, melasmas, wrinkles, warts, blemished skin, hyperpigmented skin, hyperkeratotic skin, inflammatory dermatoses, age-related skin changes and psoriasis.

59. A method for enhancing epidermal exfoliation and/or for enhancing epidermal cell renewal which comprises topically administering to a subject on an area of skin an effective amount of a composition comprising (1) an acid protease which is enzymatically active below about pH 5.5 and which is insignificantly active at or above pH 5.5 present in an amount of 4% by weight of the final composition having an activity of 4000 HUT units/mg; and (2) an acidic buffer comprising lactic acid present in an amount of 3% by weight of the final composition which temporarily lowers the surface pH of the skin to below about pH 5.5 for a period of time effective for enhancing epidermal exfoliation and/or epidermal cell renewal, the acidic buffer being subject to neutralization by natural epidermal processes, such that the surface pH of the skin returns to about pH 5.5.

60. A method for enhancing epidermal exfoliation and/or for enhancing epidermal cell renewal which comprises topically administering to a subject on an area of skin an effective amount of a composition comprising (1) an acid protease which is enzymatically active below about pH 5.5 and which is insignificantly active at or above pH 5.5 present in an amount of 4% by weight of the final composition having an activity of 4000 HUT units/mg; and (2) an acidic buffer comprising lactic acid present in an amount of 1.5% by weight of the final composition which temporarily lowers the surface pH of the skin to below about pH 5.5 for a period of time effective for enhancing epidermal exfoliation and/or epidermal cell renewal, the acidic buffer being subject to

neutralization by natural epidermal processes, such that the surface pH of the skin returns to about pH 5.5.

61. The method according to claim 60 in which the increase in exfoliation is 136% as compared to control.

62. The method according to claim 59 in which the acid protease is pepsin.

10 63. The method according to claim 60 in which the acid protease is pepsin.

64. A method for enhancing epidermal exfoliation and/or for enhancing epidermal cell renewal which comprises
15 topically administering to a subject on an area of skin an effective amount of a composition comprising (1) an acid protease which is enzymatically active below about pH 5.5 and which is insignificantly active at or above pH 5.5 present in an amount of 4% by weight of the final composition having an
20 activity of 4000 HUT units/mg; and (2) an acidic buffer comprising citric acid present in an amount of 3% by weight of the final composition which temporarily lowers the surface pH of the skin to below about pH 5.5 for a period of time effective for enhancing epidermal exfoliation and/or
25 epidermal cell renewal, the acidic buffer being subject to neutralization by natural epidermal processes, such that the surface pH of the skin returns to about pH 5.5.

65. A method for enhancing epidermal exfoliation
30 and/or for enhancing epidermal cell renewal which comprises topically administering to a subject on an area of skin an effective amount of a composition comprising (1) an acid protease which is enzymatically active below about pH 5.5 and which is insignificantly active at or above pH 5.5 present in
35 an amount of 4% by weight of the final composition having an activity of 4000 HUT units/mg; and (2) an acidic buffer comprising lactic acid present in an amount of 1.5% by weight of the final composition and citric acid present in an amount

of 1.5% by weight of the final composition which temporarily lowers the surface pH of the skin to below about pH 5.5 for a period of time effective for enhancing epidermal exfoliation and/or epidermal cell renewal, the acidic buffer being
5 subject to neutralization by natural epidermal processes, such that the surface pH of the skin returns to about pH 5.5.

66. The method according to claim 64 in which the acid protease is pepsin.

10

67. The method according to claim 65 in which the acid protease is pepsin.

15

20

25

30

35

1/3




FIG. 1

2/3




FIG. 2

3/3




FIG. 3

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/10427

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 7/48

US CL :424/401; 514/844, 845, 846, 847, 944

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/401; 514/844, 845, 846, 847, 944

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONEElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)
NONE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,431,913 A (PHILIPPS) 11 July 1995, see column 4, lines 20-60.	1-67
A	US 4,363,815 A (YU et al.) 14 December 1982, see columns 4-7.	1-67

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A document defining the general state of the art which is not considered to be of particular relevance	*X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E earlier document published on or after the international filing date	*Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z document member of the same patent family
*O document referring to an oral disclosure, use, exhibition or other means	
*P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

20 AUGUST 1997

Date of mailing of the international search report

11 SEP 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

J. VENKAT

Telephone No. (703) 308-2351